


Agro-  
Imperial Agricultural Research  
Institute, Delhi  
Biological  
& Medical  
Series  
June, 1913  
Botanical Series.  
Vol. VI, No. 1.

STORAGE

# Memoirs of the Department of Agriculture in India

NOTES ON POLLINATION AND CROSS-FERTILISA-  
TION IN THE COMMON RICE PLANT  
ORYZA SATIVA, LINN

BY  
G. P. HECTOR, M.A., B.Sc  
*Economic Botanist, Bengal*



AGRICULTURAL RESEARCH INSTITUTE, PUSA  
PUBLISHED FOR  
THE IMPERIAL DEPARTMENT OF AGRICULTURE IN INDIA  
BY  
THACKER, SPINK & CO., CALCUTTA  
W. THACKER & CO., 2, CREED LANE, LONDON

Price, Rs. 1.







THE ROYAL CANADIAN INSTITUTE  
MEMOIRS OF THE  
DEPARTMENT OF AGRICULTURE  
IN INDIA

BOTANICAL SERIES

VOL. VI



AGRICULTURAL RESEARCH INSTITUTE, PUSA

PUBLISHED FOR  
THE IMPERIAL DEPARTMENT OF AGRICULTURE IN INDIA

BY  
THACKER, SPINK & CO., CALCUTTA  
W. THACKER & CO., 2, CREED LANE, LONDON

CALCUTTA :  
PRINTED BY THACKER, SPINK AND CO.



# CONTENTS.

## VOL. VI.

|  | PAGE. |
|--|-------|
| No. 1. HECTOR, G. P. Notes on Pollination and Cross-Fertilization in the Common Rice Plant, <i>Oryza Sativa</i> , Linn. ...  | 1     |
| No. 2. SHAW, F. J. F. A Sclerotial Disease of Rice (with 3 plates—1 coloured)...   | 11    |
| No. 3. HOWARD, GABRIELLE L. C. Studies in Indian Tobaccos, No. 3. The Inheritance of Characters in <i>Nicotiana tabacum</i> , L. (with 25 plates) ...  | 25    |
| No. 4. LEAKE, H. MARTIN, and RAM PRASAD. Studies in Indian Cottons, Part I. The Vegetative Characters (with 3 maps and 19 plates—5 coloured) ...   | 115   |
| No. 5. BUTLER, E. J., and ABDUL HAFIZ KHAN. Red Rot of Sugarcane (with one plate) ...  | 151   |
| No. 6. BUTLER, E. J., and ABDUL HAFIZ KHAN. Some new Sugarcane Diseases (with 6 plates—1 coloured) ...   | 181   |
| No. 7. GRAHAM, R. J. D. Preliminary Note on the Classification of Rice in the Central Provinces (with 4 plates and one coloured map)   | 209   |
| No. 8. HOWARD, A., LEAKE, H. M., and HOWARD, GABRIELLE L. C. The Influence of the Environment on the Milling and Baking Qualities of Wheat in India. No. 3. The Experiments of 1911—12 (with one coloured plate and a map) ... | 231   |





June 1913

BOTANICAL SERIES

Vol. VI, No. 1.

# MEMOIRS OF THE DEPARTMENT OF AGRICULTURE IN INDIA

NOTES ON POLLINATION AND CROSS-FERTILISA-  
TION IN THE COMMON RICE PLANT,  
ORYZA SATIVA, LINN.

BY

G. P. HECTOR, M.A., B.SC.

*Economic Botanist, Bengal*



AGRICULTURAL RESEARCH INSTITUTE, PUSA

PUBLISHED FOR

THE IMPERIAL DEPARTMENT OF AGRICULTURE IN INDIA

BY

THACKER, SPINK & CO., CALCUTTA  
W. THACKER & CO., 2, CREED LANE, LONDON

MEMOIRS OF THE  
DEPARTMENT OF AGRICULTURE  
IN INDIA

NOTES ON POLYVALENCE AND COMPARATIVES  
TOWARDS THE COMMON HONEY BEES  
ON THE NATIVE FORM

CALCUTTA :

PRINTED BY THACKER, SPINK AND CO.

AGRICULTURAL DEPARTMENT, CALCUTTA

AGRICULTURAL DEPARTMENT, CALCUTTA

THACKER, SPINK AND CO. LTD.  
PRINTERS, CALCUTTA



## CONTENTS.

|                          |     |     |     | Page. |
|--------------------------|-----|-----|-----|-------|
| I. Introductory          | ... | ... | ... | 1     |
| II. Pollination          | ... | ... | ... | 2—4   |
| III. Cross-fertilisation | ... | ... | ... | 5—10  |





Digitized by the Internet Archive  
in 2008 with funding from  
Microsoft Corporation



# NOTES ON POLLINATION AND CROSS-FERTILISATION IN THE COMMON RICE PLANT, *ORYZA SATIVA*, LINN.

BY

G. P. HECTOR, M.A., B.SC.,

*Economic Botanist, Bengal.*

## I.—INTRODUCTORY.

DURING the past two years a large number of varieties of rice, cultivated in the districts of Lower Bengal, have been grown on the Dacca farm, with the object of studying their characters in single plant cultures and of ascertaining to what extent, if any, natural cross-fertilisation takes place, as a correct estimation of this latter point and of the precautions likely to be necessary to keep varieties pure, is an essential preliminary to the work of successfully distributing seed of improved types. This preliminary work is necessary in the case of all crops, but particularly so in the case of a crop like rice, in which the number of cultivated varieties, differing frequently by most minute points of difference, is very large, and the risk of accidental mixture of seed very great, owing to the peculiar conditions under which rice is grown.

Though the results obtained up to date are far from complete, they have shown that while some types are much superior to others and self-fertilisation is the normal process, cross-fertilisation does undoubtedly take place and under certain circumstances may considerably affect the successful introduction of an improved type into any particular locality where other inferior types are commonly grown. This question of pollination and



cross-fertilisation in rice has not been studied in detail in India, and in the following notes it is proposed to give a preliminary account of our observations on the subject, as far as the districts of Lower Bengal are concerned.

## II.—POLLINATION.

A superficial examination of the flower of rice would lead one to believe that cross-pollination by the agency of the wind was the normal method. Shortly after the opening of the glumes the anthers may be seen hanging downwards from their filaments in such a way as almost to preclude the pollen from reaching the stigma of the same flower if dehiscence has not already taken place before this pendent position is reached, and frequently also the two feathery stigmas may be seen protruding one on either side from between the open glumes. Though no work has hitherto been done in India on the subject, much has been done in other countries, and a brief summary of previous literature is given below. Knuth\* states that the species is allogamous and that the anthers are still closed when they emerge from the glumes, and do not dehisce till after they have bent over and hang downwards. More recently a Japanese worker, M. Akeipine,† has given a detailed account of the morphology of the flower and of the flowering of a race of swamp-rice, known as “Akaghé,” occurring in Northern Japan. Only a summary‡ of this paper has been seen by us. According to the summary, the author discusses in detail the climatic factors affecting the flowering and setting of the grain, describes the method of pollination, and gives cytological details regarding the fertilisation process. He gives no cases of the occurrence of cross-fertilisation, and concludes, from observations and experiments on the flower, that self-fertilisation is what usually occurs, pollination according to the writer taking place just before the flowers open. Lastly, Fruwirth and Van der

---

\* Hand-book of Flower Pollination, Vol. III, 1909, p. 521.

† “On the Flowers and Flowering of *O. sativa*,” Agric. Gazette, Nôgyô-Sekai, 1910-11.

‡ Cf. Bot. Centralblatt, No. 41 of 1911, pp. 370-71.

Stok\* summarize fully all the work done up to date, chiefly by the latter in Java, on the subject of rice-breeding in general, including the question of pollination and cross-fertilisation. According to these authorities, self-fertilisation is the usual process, but cross-fertilisation may not infrequently take place between neighbouring plants, and must be taken account of in all breeding work.

According to our observations on the Dacca farm, the actual process of pollination is as follows :—

The spikelets, or flowers, mature from above downwards in a fairly regular sequence, and in the case of any individual flower the opening of the glumes and pollination takes place, as a rule, on the same day as that on which it emerges above the level of the leaf-sheath if this happens before mid-day, or at latest in the course of the forenoon of the following day. In almost every normal case, dehiscence of the anthers and pollination take place practically simultaneously with the opening of the glumes, sometimes even before they open at all, and at the time of dehiscence the stigmas are still enclosed and the anthers still within the shelter of the glumes. Hence self-fertilisation appears to be the usual method. In the case of *Aus* varieties, which in Lower Bengal flower during the months of May and June, the opening of the glumes and dehiscence of the anthers commences usually between the hours of 7 and 8 A.M. and continues till about 10 A.M. when it stops for the day, but in the case of the *Aman* varieties which generally begin to flower in late October or early November, not till later in the forenoon, beginning usually between 9 and 10 A.M., and continuing till mid-day. The later time of day at which the flowers of the *Aman* varieties open is probably due to the much lower temperatures in October and November.

Immediately before the flowers open and dehiscence takes place it may be seen that the top of the anthers is just touching the concave roof formed by the glumes which are still closed,

---

\* Fruwirth, Die Züchtung der Landwirtschaftlichen Kulturpflanzen, B. V., 1912, pp. 36—54.



and if at this stage the glumes are gently pulled apart, the anthers may be seen to dehisce and the pollen to fall in showers on to the feathery stigmas within. It is probable that the pressure of the anthers—which at this stage are being pushed up by the rapidly elongating filaments from below—on the glumes, together with the slight shock caused by the opening of the glumes and the consequent access of air and sunlight, are the immediate causes which bring about dehiscence.

Immediately after dehiscence, the glumes diverge a little further apart, the filaments rapidly elongate till the anthers protrude about their own length above the tip of the glumes, and then bend slowly outwards and downwards so that the anthers ultimately assume a pendent position. But before this stage is reached, in every normal case the anthers have already lost almost all their pollen and pollination has taken place before they emerged at all.

Both the extent to which the flowers open and the length of time they remain open vary considerably, but we have not been able to discover any differences in varieties with regard to this. It appears to be largely due to atmospheric conditions at the time. In the warmer and moister months of May and June the flowers of the *Aus* varieties open earlier in the morning, but appear to open to a less extent, and to remain open a shorter time than in the case of *Aman* varieties which flower in the colder and drier months of October-November. According to our observations, the whole process from the time of the opening of the glumes and dehiscence of the anthers till the time when the latter assume the pendent position, occupies on an average only about fifteen minutes, in both *Aus* and *Aman* varieties, but in the case of *Aus* varieties the flowers seldom remain open for a longer period than half an hour in all, while in the case of *Aman* varieties they may remain open for periods varying from about an hour to an hour and a half. If the weather is wet and rainy at the time when the flowers should normally open, as is frequently the case when the *Aus* varieties flower, they may not open at all, or if they do, they often do not close again, and a large percentage of

such flowers set no grain. It is a matter of common observation that a far larger number of empty husks are to be found in the heads of *Aus* varieties than in *Aman* varieties, due probably to the wet weather which often occurs in Lower Bengal when the *Aus* rices commence to flower.

In about four days a whole inflorescence will ordinarily have been pollinated in the way described above.

### III.—CROSS-FERTILISATION.

Though the above is the normal process, the method of pollination by no means precludes the possibility of cross-pollination taking place. It has been pointed out above that the stigmas are not infrequently to be seen protruding from between the open glumes, while the stamens have not lost all their pollen when they assume the hanging position, so that there is every possibility, should self-pollination have failed, of cross-pollination resulting, and we have collected evidence to prove that it is by no means unknown, and under favourable weather conditions, bright sunshine with a gentle breeze, possibly takes place to a greater extent than we have as yet been able to prove. It is practically certain, however, that such cross-pollination as does take place is confined to neighbouring plants in the same plot, and that the pollen of one plant seldom if ever succeeds in reaching the stigma of a plant more than a foot or two away.

In the following paragraphs an account is given of the evidence we have so far collected as to the extent to which cross-pollination has been found to occur.

In 1911 single plant cultures of 150 types of transplanted *Aman* varieties were grown at Dacca farm from single plants selected in the season 1910. No cases of splitting crosses were found in these, and all were passed as pure at harvest time, but in four plots a few variant plants were found. Subsequent examination, however, in the laboratory, of the seed of certain white-grained varieties, proved that in some, red-grained seeds were to be found which had escaped detection at harvest time, and possibly there were others which were not discovered. Six



such cases were found, and in each case as many red-grained seeds as could be picked out were sown and the seedlings transplanted separately in 1912, every precaution being taken to prevent accidental mixture. In three of the six plots the red seeds have given rise to a few white-grained plants, proving that these were heterozygous as regards seed colour.

To return to the variant plants mentioned above, seed of four of these, which it was suspected might be crosses, was carefully preserved, sown and transplanted separately in 1912, and all have given rise to a variety of types as described below :—

The first of these (labelled Ba 1), was found in a variety named Gobrabali, a coarse rice from Bakarganj, and differed from the type chiefly in having reddish-coloured awns. This in 1912 split into four types—

- |   |     |    |
|---|-----|----|
| (1) Red awns, two inner glumes at flowering time pale green with red tip, turning yellow when ripe, stigmas black | ... | 60 |
| (2) Red awns, inner glumes at flowering time dark green with red tip, turning mottled brown, stigmas black        | ... | 13 |
| (3) White awns, inner glumes at flowering time pale green with no tip, turning yellow, stigmas white              | ... | 19 |
| (4) White awns, inner glumes at flowering time dark green with no tip, turning brown, stigmas white               | ... | 4  |

Ratio red awns to white awns, 3:17 : 1

The second (labelled R 20) was found in a variety named Ukulmadhu from Rajshahi. The parent plant had a small, fine grain, the two small outer glumes yellow, and the inner glumes yellow with brownish coloured apex. This in 1912 split into three main types—

- |  |     |     |     |          |
|--|-----|-----|-----|----------|
| (1) Small outer glumes yellow, inner glumes at first green with reddish tip, turning yellow with reddish tip, stigmas purple | ... | ... | ... | 5 noted. |
| (2) Ditto, but with stigmas white  | ... | ... | ... | 5 "      |
| (3) Small outer glumes coloured reddish-brown, inner glumes at first reddish with no tip, turning black, stigmas white       | ... | ... | ... | 2        |

The only obvious difference between types (1) and (2) in this case lay in the colour of the stigma which can be accurately determined only before pollination has taken place. Unfortunately

the plot was not examined in detail till after flowering was almost over, and the colour of the stigma could be determined in only a small number of plants.

The third and fourth examples will be treated together. These were red-grained plants picked from white-grained plots, the red-grained plants in both cases being otherwise almost indistinguishable from the remaining white-grained plants in the plots. The first of these (labelled M 1) was found in a variety named Aman paddy from Mymensing, and the second (labelled B 15) in a variety named Bankalam from Bogra. These both split in 1912 into red and white-grained plants, otherwise almost indistinguishable except for grain colour, in the proportions given below—

|                           |                       |         |          |
|---------------------------|-----------------------|---------|----------|
| 1. M1, red-grained, 1911  | ... 1912, red-grained | ... 145 | } ratio. |
|                           | white-grained         | ... 55  |          |
| 2. B15, red-grained, 1911 | ... 1912, red-grained | ... 132 | } ratio. |
|                           | white-grained         | ... 68  |          |

Besides these four cases, much further evidence has been afforded by a series of Dinajpur varieties started from single plants selected in 1911. The seed of these varieties was collected from Dinajpur, partly from cultivators and partly from the Bazaars, and all were very mixed. In 1911, eighty-six single plants were selected from these Dinajpur plots and these were sown and transplanted separately in 1912 and no fewer than seven, or a percentage of 8·15, have given rise to a mixture of types. These are described below in detail.

In the first (labelled D 28), the parent plant had the small outer glumes coloured reddish-brown, and the inner glumes yellow with a dark apex. This in 1912 split into the following four main types :—

- |  |   |        |
|--|---|--------|
| (1) Outer glumes coloured reddish-brown, inner glumes at flowering time green with red tip, turning yellow with dark tip, stigmas black... | 2 | noted. |
| (2) Ditto with stigmas white   | 6 | "      |
| (3) Taller, more robust plant, outer glumes coloured reddish-brown, inner glumes at flowering time reddish, turning black, stigmas black   | 2 |        |
| (4) Ditto with stigmas white   | 1 |        |



As in the case of R 20 described above, here again it was too late to determine accurately the stigma colour in types (1) and (2), the only obvious point of difference between them, when the plot was examined, except in the cases noted.

In the second (labelled D 6), the parent plant had a red grain and in 1912 split into four main types, two with red and two with white grains, as noted below.

| D 6—Red-grained, 1911.   |                               |  |                                |
|--|-------------------------------|--|--------------------------------|
| (1) Reddish leaf-sheaths and stem above nodes, grain white ... 14. | (2) Ditto, grain amber ... 1. | (3) Green leaf-sheaths and stems, grain red of various shades ... 43 | (4) Ditto, grain white ... 26. |

In the third (labelled D 7), the parent was red-grained, and split into three main types, two with red grains and one with white, in the proportions given—

|                                       |     |     |        |
|---------------------------------------|-----|-----|--------|
| 1. Glumes yellow, grain red           | ... | ... | ... 50 |
| 2. Glumes mottled brown, grain red... | ... | ... | ... 34 |
| 3. Glumes mottled brown, grain white  | ... | ... | ... 13 |

The remaining four Dinajpur examples are similar to those of M 1 and B 15 described above, *viz.*, red-grained parent plants splitting into red and white-grained offspring, almost indistinguishable except for the colour of the grain. The details of these four cases are as follows :—

| PARENT 1911. |              | OFFSPRING 1912. |                | RATIO R. : W. |
|--------------|--------------|-----------------|----------------|---------------|
|              | Red-grained. | Red-grained.    | White-grained. |               |
| D 17         | ...          | 72              | 21             | 3.4 : 1       |
| D 18         | ...          | 72              | 24             | 3 : 1         |
| D 35         | ...          | 51              | 21             | 2.42 : 1      |
| D 38         | ...          | 68              | 32             | 2.12 : 1      |

From a perusal of the above figures, together with those of M 1 and B 15 quoted above, it would seem that as regards colour of grain segregation is taking place in the simple Mendelian ratio of 3 : 1.\* Cross-fertilisation experiments are in

\* Cf. Van der Stok, l. c., p. 49.

progress with the object of verifying this and also of determining the mode of inheritance of certain other characters.

In addition to the examples described above of undoubted splitting crosses, stray variant plants have also been found in twelve of the 1912 pure line plots started from single plants selected in 1910, and in fifteen of the 1912 Dinajpur series started from single plants selected in 1911, amounting to about fifty in all. Some of these may prove to be accidental mixtures but some cannot be matched with any of our other types, and have all the appearance of being the  $F_1$  generation of crosses which must have taken place within our area in 1911. Seed of all these is being preserved and will be sown in 1913 and the results noted.

From the above cases there is reason to believe that natural crossing in rice is more common than was at first supposed. Moreover, under the conditions in which rice is grown by the cultivator, whose varieties are never free from mixtures, it would probably take place to a considerably greater extent than in our area at Dacca. Even if it did not occur to any greater extent than the cases cited would indicate, it would be quite sufficient to account for the extraordinarily large number of types which are to be found, when one takes into consideration the extent and great antiquity of the cultivation. Certain facts also which have been brought to our notice lend further evidence to this belief. For example, a cultivator from the neighbourhood of Chandpur recently informed us that on his own land within the past twelve years the number of distinct types to be found in his fields has increased from eight or ten to almost a hundred, although he has imported no new varieties.

The main conclusions to be drawn from the above results are :—

1. That in Lower Bengal under favourable conditions cross-fertilisation may take place in rice to an extent which may be provisionally estimated at about 4 per cent.

2. That this cross-fertilisation takes place wholly through the agency of the wind and would seem to be effective only between flowers of adjacent plants to a radius of a few feet.



3. That as regards certain characters at least, *e.g.*, grain colour, segregation along Mendelian lines appears to take place.

4. That so long as seed of a variety is kept free from accidental mixture there is no risk of contamination from cross-fertilisation, but that if seed gets mixed, cross-fertilisation will undoubtedly take place between adjacent plants in a plot and to an extent sufficient in a few years' time to reduce a variety to a number of splitting types. Hence the imperative necessity of taking every precaution to keep seed of varieties free from accidental mixtures.

Dacca, }  
10th January 1913. }

# PUBLICATIONS OF THE IMPERIAL DEPARTMENT OF AGRICULTURE IN INDIA.

[TO BE HAD FROM MESSRS. THACKER, SPINK & CO., CALCUTTA.]

- Annual Report of the Imperial Department of Agriculture in India, for the year 1904-05. Price, As. 12 or 1s. 2d. (*Out of print.*)
- Report of the Imperial Department of Agriculture in India, for the years 1905-06 and 1906-07. Price, As. 6 or 7d.
- Report of the Agricultural Research Institute and College, Pusa (including Report of the Imperial Cotton Specialist), for the years 1907-09. Price, As. 4.
- Report of the Agricultural Research Institute and College, Pusa (including Report of the Imperial Cotton Specialist), for the year 1909-10. Price, As. 4 or 5d.
- Report of the Agricultural Research Institute and College, Pusa (including Report of the Imperial Cotton Specialist), for 1910-11. Price, As. 6 or 7d.
- Report of the Agricultural Research Institute and College, Pusa (including Report of the Imperial Cotton Specialist), for 1911-12. Price, As. 6 or 7d.
- Report on the Progress of Agriculture in India, for the years 1907-09. Price, As. 6 or 7d.
- Report on the Progress of Agriculture in India, for the year 1909-10. Price, As. 6 or 7d.
- Report on the Progress of Agriculture in India, for 1910-11. Price, As. 12 or 1s. 3d.
- Report on the Progress of Agriculture in India, for 1911-12. Price, As. 6 or 7d.
- Proceedings of the Board of Agriculture in India, held at Pusa on the 6th January 1905 and following days (with Appendices). Price, As. 8 or 9d.
- Proceedings of the Board of Agriculture in India, held at Pusa on the 15th January 1906 and following days (with Appendices). Price, As. 12 or 1s. 2d.
- Proceedings of the Board of Agriculture in India, held at Cawnpur on the 18th February 1907 and following days (with Appendices). Price, Re. 1-2 or 1s. 6d.
- Proceedings of the Board of Agriculture in India, held at Pusa on the 17th February 1908 and following days (with Appendices). Price, As. 8 or 9d.
- Proceedings of the Board of Agriculture in India, held at Nagpur on the 15th February 1909 and following days (with Appendices). Price, As. 8 or 9d.
- Proceedings of the Board of Agriculture in India, held at Pusa on the 21st February 1910 and following days (with Appendices). Price, As. 8 or 9d.
- Proceedings of the Board of Agriculture in India, held at Pusa on the 20th November 1911 and following days (with Appendices). Price, As. 10 or 1s.
- Standard Curriculum for Provincial Agricultural Colleges as recommended by the Board of Agriculture, 1908. Price, As. 4 or 5d.
- The "*Agricultural Journal of India.*" A Quarterly Journal dealing with subjects connected with field and garden crops, economic plants and fruits, soils, manures, methods of cultivation, irrigation, climatic conditions, insect pests, fungus diseases, co-operative credit, agricultural cattle, farm implements and other agricultural matters in India. Illustrations, including coloured plates, form a prominent feature of the Journal. It is edited by the Agricultural Adviser to the Government of India, assisted by an Advisory Committee of the Staff of the Agricultural Research Institute, Pusa. *Annual Subscription*, Rs. 6 or 8s. Single copy, Rs. 2.



MEMOIRS OF THE DEPARTMENT OF AGRICULTURE IN INDIA are issued from time to time as matter is available, in separate series, such as Chemistry, Botany, Entomology and the like.

### BOTANICAL SERIES.

- Vol. I, No. I. Studies in Root Parasitism. The Haustorium of *Santalum album*.—Part I.—Early Stages by C. A. BARBER, M.A., F.L.S. Price, Re. 1.  
Part II.—The Structure of the Mature Haustorium and the Inter-relations between Host and Parasite by C. A. BARBER, M.A., F.L.S. Price, Rs. 3.
- Vol. I, No. II. Indian Wheat Rusts by E. J. BUTLER, M.B., F.L.S. ; and J. M. HAYMAN, D.V.S. Price, Rs. 3.
- Vol. I, No. III. Fungus Diseases of Sugarcane in Bengal by E. J. BUTLER, M.B., F.L.S. Price, Rs. 3.
- Vol. I, No. IV. *Gossypium obtusifolium*, Roxburgh, by I. H. BURKILL, M.A. Price, Re. 1.
- Vol. I, No. V. An Account of the Genus *Pythium* and some *Chytridiaceæ* by E. J. BUTLER, M.B., F.L.S. Price, Rs. 4-8.
- Vol. I, No. VI. *Cephaleuros virescens*, Kunze ; The Red Rust of Tea by HAROLD H. MANN, D.Sc. ; and C. M. HUTCHINSON, B.A. Price, Rs. 4.
- Vol. II, No. I. Some Diseases of Cereals caused by *Sclerospora graminicola* by E. J. BUTLER, M.B., F.L.S. Price, Re. 1-8.
- Vol. II, No. II. The Indian Cottons by G. A. GAMMIE, F.L.S. Price, Rs. 7-8.
- Vol. II, No. III. Note on a Toxic Substance excreted by the Roots of Plants by F. FLETCHER, M.A., B.Sc. Price, Re. 1-8.
- Vol. II, No. IV. Studies in Root Parasitism III. The Haustorium of *Olaæ scandens* by C. A. BARBER, M.A., F.L.S. Price, Rs. 2-8.
- Vol. II, No. V. Studies in Root Parasitism IV. The Haustorium of *Canjiera Rheedii* by C. A. BARBER, M.A., F.L.S. Price, Rs. 2-8.
- Vol. II, No. VI. Some Experiments in the Hybridising of Indian Cottons by P. F. Fyson, B.A., F.L.S. Price, Re. 1-8.
- Vol. II, No. VII. The Varietal Characters of Indian Wheats by ALBERT HOWARD, M.A., A.R.C.S., F.L.S. ; and GABRIELLE L. C. HOWARD, M.A. Price, Re. 1.
- Vol. II, No. VIII. The Mulberry Disease caused by *Coryneum mori*, Nom. in Kashmir, with notes on other Mulberry Diseases, by E. J. BUTLER, M.B., F.L.S. Price, Re. 1-8.
- Vol. II, No. IX. The Wilt Disease of Pigeon-Pea and the Parasitism of *Neocosmospora vasinfecta*, Smith, by E. J. BUTLER, M.B., F.L.S. Price, Rs. 3.
- Vol. III, No. I. Studies in Indian Tobaccos, No. 1. The Types of *Nicotiana rustica*, L., Yellow Flowered Tobacco, by ALBERT HOWARD, M.A., A.R.C.S., F.L.S. ; and GABRIELLE L. C. HOWARD, M.A. Price, Rs. 4.
- Vol. III, No. II. Studies in Indian Tobaccos, No. 2. The Types of *Nicotiana tabacum*, L., by ALBERT HOWARD, M.A., A.R.C.S., F.L.S. ; and GABRIELLE L. C. HOWARD, M.A. Price, Rs. 9.
- Vol. III, No. III. Studies in Indian Fibre Plants, No. 1. On two varieties of *Sann, Crotalaria juncea*, L., by ALBERT HOWARD, M.A., A.R.C.S., F.L.S. ; and GABRIELLE L. C. HOWARD, M.A. Price, Re. 1.
- Vol. III, No. IV. The Influence of the Environment on the Milling and Baking Qualities of Wheat in India. No. 1.—The Experiments of 1907-08 and 1908-09. By ALBERT HOWARD, M.A., A.R.C.S., F.L.S. ; H. M. LEAKE, M.A., F.L.S. ; and GABRIELLE L. C. HOWARD, M.A. Price, Re. 1-8.
- Vol. III, No. V. The Bud-Rot of Palms in India by E. J. BUTLER, M.B., F.L.S. Price, Rs. 2.
- Vol. III, No. VI. The Economic Significance of Natural Cross-fertilisation in India by ALBERT HOWARD, M.A., A.R.C.S., F.L.S. ; GABRIELLE L. C. HOWARD, M.A. ; and ABDUR RAHMAN KHAN. Price, Rs. 4-8.
- Vol. IV, No. I. Millets of the Genus *Setaria* in the Bombay Presidency and Sind by G. A. GAMMIE, F.L.S., Imperial Cotton Specialist. Price, Re. 1.

## BOTANICAL SERIES—contd.

- Vol. IV, No. II. Studies in Indian Fibre Plants. No. 2.—On Some New Varieties of *Hibiscus cannabinus*, L., and *Hibiscus Sabdarifa*, L., by ALBERT HOWARD, M.A., A.R.C.S., F.L.S.; and GABRIELLE L. C. HOWARD, M.A. Price, Rs. 3.
- Vol. IV, No. III. Notes on the Incidence and Effect of Sterility and Cross-fertilisation in the Indian Cottons by H. M. LEAKE, M.A. (Cantab.), F.L.S.; and RAM PRASAD. Price, Re. 1.
- Vol. IV, No. IV. Note on the Inheritance of Red Colour and the Regularity of Self-fertilisation in *Cochorus capsularis*, the common Jute Plant, by I. H. BURKILL, M.A.; and R. S. FINLOW, B.Sc., F.C.S. Price, Re. 1.
- Vol. IV, No. V. Observations on Certain Extra-Indian Asiatic Cottons by H. M. LEAKE, M.A., F.L.S.; and RAM PERSHAD, Asst. to the Economic Botanist, U. P. Price, Re. 1-8.
- Vol. IV, No. VI. The Morphology and Parasitism of *Rhizoctonia* by F. J. F. SHAW, B.Sc., A.R.C.S., F.L.S. Price, Rs. 2.
- Vol. V, No. I. The Inheritance of some Characters in Wheat—I, by A. HOWARD, M.A., A.R.C.S., F.L.S.; and GABRIELLE L. C. HOWARD, M.A. Price, Re. 1.
- Vol. V, No. II. The Influence of the Environment on the Milling and Baking Qualities of Indian Wheats. No. 2.—The Experiments of 1909-10 and 1910-11, by A. HOWARD, M.A., A.R.C.S., F.L.S.; GABRIELLE L. C. HOWARD, M.A.; and H. M. LEAKE, M.A., F.L.S. Price, Re. 1.
- Vol. V, No. III. The Varieties of Soy Beans found in Bengal, Bihar and Orissa and their Commercial possibilities, by E. J. WOODHOUSE, M.A.; and C. S. TAYLOR, B.A. Price, Rs. 2.
- Vol. V, No. IV. On *Phytophthora parasitica* nov. spec. by J. F. DASTUR, B.Sc. (*In the press.*)
- Vol. V, No. V. Studies in *Peronosporaceæ* by E. J. BUTLER, M.B., F.L.S., Imperial Mycologist, and G. S. KULKARNI, L.Ag., Mycological Assistant, Bombay. Price, Rs. 2.
- Vol. VI, No. I. Notes on Pollination and Cross-Fertilisation in the Common Rice Plant, *Oryza sativa*, Linn., by G. P. HECTOR, M.A., B.Sc. Price, Re. 1.
- Vol. VI, No. II. A Sclerotial Disease of Rice by F. J. F. SHAW, B.Sc. (Lond.), A.R.C.S., F.L.S. (*In the press.*)

## CHEMICAL SERIES.

- Vol. I, No. I. The Composition of Indian Rain and Dew by J. WALTER LEATHER, Ph.D., F.I.C. Price, Re. 1.
- Vol. I, No. II. The Composition of the Oil Seeds of India by J. W. LEATHER, Ph.D., F.I.C. Price, Re. 1.
- Vol. I, No. III. The Pot-Culture House at the Agricultural Research Institute, Pusa, by J. W. LEATHER, Ph.D., F.I.C. Price, Rs. 3.
- Vol. I, No. IV. Experiments on the Availability of Phosphates and Potash in Soils by J. W. LEATHER, Ph.D., F.I.C. Price, Re. 1-8.
- Vol. I, No. V. The Construction of Drain Gauges at Pusa by M. H. ARNOTT, M.INST.C.E., with a Preface by J. W. LEATHER, Ph.D., F.I.C. Price, Rs. 3.
- Vol. I, No. VI. The Loss of Water from Soil during Dry Weather by J. WALTER LEATHER, Ph.D., F.I.C. Price, Rs. 2.
- Vol. I, No. VII. The System Water, Calcium Carbonate, Carbonic Acid by J. WALTER LEATHER, Ph.D., F.I.C.; and JATINDRA NATH SEN, M.A., F.C.S. Price, Re. 1.
- Vol. I, No. VIII. Water Requirements of Crops in India by J. WALTER LEATHER, Ph.D., F.I.C. Price, Rs. 3.
- Vol. I, No. IX. The Nature of the Colour of Black Cotton Soil by H. E. ANNETT, B.Sc. (Lond.), F.C.S., M.S.E.A.C. Price, Re. 1.
- Vol. I, No. X. Water Requirements of Crops in India—II, by J. WALTER LEATHER, Ph.D., F.I.C., Imperial Agricultural Chemist. Price, Rs. 2-8.
- Vol. II, No. I. The Composition of the Milk of some Breeds of Indian Cows and Buffaloes and its Variations, Part I, by A. A. MEGGITT, B.Sc. (Lond.), and H. H. MANN, D.Sc. Price, Re. 1-8.
- Vol. II, No. II. Records of Drainage in India by J. W. LEATHER, Ph.D., F.I.C. Price, Re. 1.
- Vol. II, No. III. The Rab System of Rice Cultivation in Western India by H. H. MANN, D.Sc.; N. V. JOSHI, B.A., B.Sc., L.Ag.; and N. V. KANITKAR, B.Ag. Price, Re. 1.



## CHEMICAL SERIES—*contd.*

- Vol. II, No. IV. The Composition of the Milk of some Breeds of Indian Cows and Buffaloes and its Variations, Part II, by A. A. MEGGITT, B.Sc. ; and H. H. MANN, D.Sc. Price, Re. 1-8.
- Vol. II, No. V. A contribution to the knowledge of the Black Cotton Soils of India by W. H. HARRISON, M.Sc., and M. R. RAMASWAMI SIVAN, B.A. Price, Re. 1.
- Vol. II, No. VI. The Date Sugar Industry in Bengal. An Investigation into its Chemistry and Agriculture. By H. E. ANNETT, B.Sc., F.C.S., M.S.E.A.C., assisted by G. K. LELE ; and BHAILAL M. AMIN. Price, Rs. 3.
- Vol. III, No. I. Evaporation from a plain Water Surface by J. W. LEATHER, Ph.D., F.I.C. (*In the press.*)
- Vol. III, No. II. Studies in the Chemistry and Physiology of the Leaves of the Betel Vine by H. H. MANN, D.Sc. ; D. L. SAHASRABUDHE and V. G. PATWARDHAN. (*In the press.*)

## ENTOMOLOGICAL SERIES.

- Vol. I, No. I. The Bombay Locust by H. M. LEFROY, M.A., F.E.S., F.Z.S. Price, Rs. 2-8.
- Vol. I, No. II. The more Important Insects injurious to Indian Agriculture by H. M. LEFROY, M.A., F.E.S., F.Z.S. Price, Rs. 3. (*Out of print.*)
- Vol. I, No. III. The Indian Surface Caterpillars of the Genus *Agrotis* by H. M. LEFROY, M.A., F.E.S., F.Z.S. ; and C. C. GHOSH, B.A. Price, Re. 1-8.
- Vol. I, No. IV. Individual and Seasonal Variations in *Helopeltis theivora*, Waterhouse, with description of a new species of *Helopeltis* by HAROLD H. MANN, D.Sc. Price, Re. 1-8.
- Vol. I, No. V. The Coccidæ attacking the Tea Plant in India and Ceylon by E. E. GREEN, F.E.S. ; and HAROLD H. MANN, D.Sc. Price, Re. 1.
- Vol. I, No. VI. The Mustard Sawfly by H. M. LEFROY, M.A., F.E.S., F.Z.S. ; and C. C. GHOSH, B.A. Price, Re. 1.
- Vol. II, No. I. The Rice Bug by H. M. LEFROY, M.A., F.E.S., F.Z.S. Price, Re. 1.
- Vol. II, No. II. Remarks on Indian Scale Insects (*Coccidæ*) by E. E. GREEN, F.E.S., F.Z.S. Price, Re. 1-8.
- Vol. II, No. III. The Red Cotton Bug by H. M. LEFROY, M.A., F.E.S., F.Z.S. Price, Re. 1.
- Vol. II, No. IV. The Castor Semi-Looper by H. M. LEFROY, M.A., F.E.S., F.Z.S. Price, Rs. 2.
- Vol. II, No. V. The Tobacco Caterpillar by H. M. LEFROY, M.A., F.E.S., F.Z.S. Price, Re. 1-8.
- Vol. II, No. VI. The Cotton Leaf-Roller by H. M. LEFROY, M.A., F.E.S., F.Z.S. Price, Re. 1-8.
- Vol. II, No. VII. Notes on Indian Scale Insects (*Coccidæ*) by H. MAXWELL-LEFROY, M.A., F.E.S., F.Z.S. Price, Re. 1-8.
- Vol. II, No. VIII. Life Histories of Indian Insects (*Coleoptera*—I), by H. MAXWELL-LEFROY, M.A., F.E.S., F.Z.S. Price, Rs. 2.
- Vol. II, No. IX. Life Histories of Indian Insects—II. Some Aquatic *Rhynchota* and *Coleoptera*, by D. NOWROJEE, B.A., Asst. to the Imperial Entomologist. Price, Re. 1.
- Vol. II, No. X. Life Histories of Indian Insects—III. The Rhinoceros Beetle (*Oryctes Rhinoceros*) and the Red or Palm Weevil (*Rhynchophorus Ferrugineus*) by C. C. GHOSH, B.A., Asst. to the Imperial Entomologist. Price, Rs. 2.
- Vol. III. The Food of Birds in India by C. W. MASON, M.S.E.A.C., edited by H. MAXWELL-LEFROY, M.A., F.E.S., F.Z.S. Price, Rs. 7-8.
- Vol. IV, No. I. Eri Silk, by H. M. LEFROY, M.A., F.E.S., F.Z.S., Imperial Entomologist ; and C. C. GHOSH, B.A., Assistant to the Imperial Entomologist. Price, Rs. 3.
- Vol. IV, No. II. Tetriginæ (Acridinæ) in the Agricultural Research Institute, Pusa, with descriptions of new species, by J. L. HANCOCK, F.Z.S. Price, Re. 1.
- Vol. IV, No. III. The Big Brown Cricket (*Brachytrypes achatinus*, Stoll), by C. C. GHOSH, B.A., Assistant to the Imperial Entomologist. Price, Re. 1.
- Vol. IV, No. IV. Life Histories of Indian Insects (Hymenoptera), by G. R. DUTT, B.A., Assistant to the Imperial Entomologist. Price, Rs. 2.
- Vol. IV, No. V. Inquiry into the Insecticidal Action of some Mineral and other compounds on Caterpillars, by H. MAXWELL-LEFROY, M.A., F.E.S., F.Z.S., Imperial Entomologist, and R. S. FINLOW, B.Sc., F.C.S., Fibre Expert to the Govt. of Bengal. Price, Re. 1-8.



## BACTERIOLOGICAL SERIES.

- Vol. I, No. I. Studies in the Bacteriological Analysis of Indian Soils, No. I. By C. M. HUTCHINSON, B.A., Imperial Agri. Bacteriologist. Price, Rs. 28.

## VETERINARY SERIES.

- Vol. I, No. I. Anaphylaxis in the larger animals, by Major J. D. E. HOLMES, D.Sc., M.A., I.C.V.D., Imperial Bacteriologist, Muktesar. Price, Rs. 2
- Vol. I, No. II. Salvarsan in the treatment of Surra in horses, dogs and rabbits, by Major J. D. E. HOLMES, D.Sc., M.A., I.C.V.D., Imperial Bacteriologist, Muktesar. Price, Re. 1-4.
- Vol. I, No. III. Some more successful Experiments on the treatment of Surra in the Camel with recommendations for Systemetic treatment, by A. S. LEESE, M.R.C.V.S. Price, Re. 1.

## BULLETINS ISSUED BY THE AGRICULTURAL RESEARCH INSTITUTE, PUSA.

- No. 1. Notes on Cotton in Behar in 1904, by H. M. LEFROY, M.A., F.E.S., F.Z.S., Imperial Entomologist. Price, As. 4 or 6d.
- No. 2. An Outbreak of Cotton Pests in the Punjab, 1905, by H. M. LEFROY, M.A., F.E.S., F.Z.S., Imperial Entomologist. Price, As. 4 or 6d.
- No. 3. The Extension of Jute Cultivation in India, by R. S. FINLOW, B.Sc., F.C.S., Jute Specialist to the Government of Eastern Bengal and Assam. Price, As. 12 or 1s. 2d. (*Out of print.*)
- No. 4. First Report on the Fruit Experiments at Pusa, by A. HOWARD, M.A. (Cantab.), A.R.C.S. (Lond.), F.L.S., Imperial Economic Botanist. Price, As. 6 or 6d.
- No. 5. Report on trials of the South African Locust Fungus in India, by E. J. BUTLER, M.B., F.L.S., Imperial Mycologist; and H. M. LEFROY, M.A., F.E.S., F.Z.S., Imperial Entomologist. Price, As. 2 or 3d.
- No. 6. The Ticks Infesting Domesticated Animals in India, by C. WARBURTON, M.A., Zoologist to the Royal Agricultural Society of England. Price, As. 4 or 6d.
- No. 7. A Preliminary Account of the Biting Flies of India, by H. M. LEFROY, M.A., F.E.S., F.Z.S., Imperial Entomologist. Price, Re. 1 or 1s. 6d. (*Out of print.*)
- No. 8. Official and Recommended Methods for use in Chemical Laboratories of the Departments of Agriculture in India, by J. WALTER LEATHER, Ph.D., F.I.C., Imperial Agricultural Chemist. Price, As. 4 or 6d.
- No. 9. Report on Cocoon Palm Disease in Travancore, by E. J. BUTLER, M.B., F.L.S., Imperial Mycologist. Price, As. 6 or 6d.
- No. 10. Treatment and Observation of Crop Pests on the Pusa Farm, by H. M. LEFROY, M.A., F.E.S., F.Z.S., Imperial Entomologist; and C. S. MISRA, B.A. Price, As. 6 or 7d.
- No. 11. On Flax Dodder, by A. HOWARD, M.A., A.R.C.S., F.L.S., Imperial Economic Botanist. Price, As. 4 or 6d.
- No. 12. The Making and Care of Lawns in India, by A. HOWARD, M.A., A.R.C.S., F.L.S., Imperial Economic Botanist. Price, As. 4 or 6d.
- No. 13. Sugarcane at the Partabgarh Experiment Station, by G. CLARKE, F.I.C., Agricultural Chemist, United Provinces; and Khan Bahadur S. M. HADI, M.R.A.C., M.R.A.S., Assistant Director of Agriculture, United Provinces. Price, As. 6 or 6d.
- No. 14. The Milling and Baking Qualities of Indian Wheats, by A. HOWARD, M.A., A.R.C.S., F.L.S.; and GABRIELLE L. C. HOWARD, M.A. Price, As. 4 or 6d.
- No. 15. Note on the Extension of Cultivation of Fibre Plants in India. Price, As. 6 or 8d.
- No. 16. Second Report on the Fruit Experiments at Pusa, by A. HOWARD, M.A. (Cantab.), A.R.C.S. (Lond.), F.L.S., Imperial Economic Botanist. Price, As. 6 or 8d.
- No. 17. The Milling and Baking Qualities of Indian Wheats, No. 2, by A. HOWARD, M.A. (Cantab.), A.R.C.S. (Lond.), F.L.S., Imperial Economic Botanist; and GABRIELLE L. C. HOWARD, M.A., Associate and late Fellow of Newnham College, Cambridge. Price, As. 6 or 8d.
- No. 18. Report on the Outbreak of Blister-Blight on Tea in the Darjeeling District in 1908-09, by W. MCRAE, M.A., B.Sc. Price, Re. 1 or 1s. 6d.
- No. 19. List of Names used in India for Common Insects, compiled in the Laboratory of the Imperial Entomologist, Pusa. Price, As. 12 or 1s. 2d.
- No. 20. Memorandum on Indian Wheat for the British Market, by Sir JAMES WILSON, K.C.S.I. Price, As. 4 or 6d.



# **BULLETINS ISSUED BY THE AGRICULTURAL RESEARCH INSTITUTE, PUSA—*contd.***

- No. 21. Memorandum regarding Leading Eucalypts suitable for India, by F. BOOTH-TUCKER, Commissioner, Salvation Army, Simla. Price, As. 4 or 5*d.*
- No. 22. The Milling and Baking Qualities of Indian Wheats, No. 3. "Some New Pusa Hybrids tested in 1910", by A. HOWARD, M.A., A.R.C.S., F.L.S., Imperial Economic Botanist; and GABRIELLE L. C. HOWARD, M.A., Personal Asst. to the Imperial Economic Botanist. Price, As. 7 or 8*d.*
- No. 23. Insecticides-Mixtures and Recipes for use against Insects in the Field, the Orchard, the Garden, and the House, by H. MAXWELL-LEFROY, M.A., F.E.S., F.Z.S., Imperial Entomologist. Second Edition, Revised and Enlarged by T. BAINBRIGGE FLETCHER, R.N., F.E.S., F.Z.S. Price, As. 12 or 1*s.* 2*d.*
- No. 24. The Indian Saltpetre Industry, by J. WALTER LEATHER, Ph.D., F.I.C., Imperial Agricultural Chemist; and JATINDRA NATH MUKHERJI, B.A., B.Sc., Second Assistant to the Imperial Agricultural Chemist. Price, As. 8 or 9*d.*
- No. 25. Report on the Flax Experiments conducted at Dooriah in 1910-11, by E. M. VANDERKERKHOVE, Flax Expert to the Bihar Planters' Association. Price, As. 6 or 7*d.*
- No. 26. Note on the present position of Cotton Investigation in India, by BERNARD COVENTRY, Offg. Inspector-General of Agriculture in India. Price, As. 2 or 3*d.*
- No. 27. Experiments on the Cultivation of Sugarcane at the Partabgarh Experimental Station, 1909-11, by G. CLARKE, F.I.C.; H. E. ANNETT, B.Sc.; and SYED ZAMIN HUSAIN, B.A. Price, As. 5 or 6*d.*
- No. 28. The Cultivation of Lac in the Plains of India, by C. S. MISRA, B.A., First Assistant to the Imperial Entomologist. Price, As. 8 or 9*d.*
- No. 29. Directions for the Cultivation of Eri Silk. Price, As. 3 or 4*d.*
- No. 30. Report on the Flax Experiments conducted at Dooriah in 1911-12, by E. M. VANDERKERKHOVE, Fibre Expert to the Bihar Planters' Association. Price, As. 1-6 or 2*d.*
- No. 31. Wheat Experiments on the Botanical Area, Cawnpur, and their bearing on Wheat Cultivation in the United Provinces, by H. MARTIN LEAKE, M.A., Economic Botanist to the Government of United Provinces, and RAMPRASAD, Assistant to the Economic Botanist. Price, As. 3 or 4*d.*
- No. 32. A Note on some Interesting Results following the Internal Administration of Arsenic in Canker and other Diseases of the foot in Horses, by J. D. E. HOLMES, M.A., D.Sc., M.R.C.V.S., Imperial Bacteriologist, Muktesar. Price, As. 2.
- No. 33. Some Aspects of the Agricultural Development of Bihar, by A. HOWARD, M.A., A.R.C.S., F.L.S., Imperial Economic Botanist. Price, As. 4 or 5*d.*

## **BOOKS.**

- "Indian Insect Pests", by H. M. LEFROY, M.A., F.E.S., F.Z.S. Price, Re. 1-8. (*Out of print.*)
- "Indian Insect Life", by H. M. LEFROY, M.A., F.E.S., F.Z.S.; and F. M. HOWLETT, B.A. 786 pp. Price, Rs. 20.
- "Wheat in India", by A. HOWARD, M.A., A.R.C.S., F.L.S.; and GABRIELLE L. C. HOWARD, M.A. 288 pp. Price, Rs. 5.







